

COMPOSITE DIFFERENTIABLE FUNCTIONS

EDWARD BIERSTONE, PIERRE D. MILMAN AND WIESŁAW PAWŁUCKI

1. INTRODUCTION

In the early 1940's, Whitney proved that every $\mathcal{C}^\infty$ even function $f(x)$ can be written $f(x) = g(x^2)$, where g is $\mathcal{C}^\infty$ [28]. About twenty years later, Glaeser (answering a question posed by Thom in connection with the $\mathcal{C}^\infty$ preparation theorem) showed that a $\mathcal{C}^\infty$ function $f(x) = f(x_1, \dots, x_m)$ which is invariant under permutations of the coordinates can be expressed $f(x) = g(\sigma_1(x), \dots, \sigma_m(x))$, where g is $\mathcal{C}^\infty$ and the $\sigma_i(x)$ are the elementary symmetric polynomials [10]. Of course, not every $\mathcal{C}^\infty$ function $f(x) = f(x_1, \dots, x_m)$ which is constant on the fibres of a (proper or semiproper) real analytic mapping $y = \varphi(x)$, $y = (y_1, \dots, y_n)$, can be expressed as a composite $f = g \circ \varphi$, where g is $\mathcal{C}^\infty$. We will say that φ has the $\mathcal{C}^\infty$ composite function property if every $\mathcal{C}^\infty$ function $f(x)$ which is “formally a composite with φ ” (see Definition 1.1 below) can be written $f = g \circ \varphi$, where $g(y)$ is $\mathcal{C}^\infty$. The theorem of Glaeser asserts that a semiproper real analytic mapping φ which is generically a submersion has the $\mathcal{C}^\infty$ composite function property. The $\mathcal{C}^\infty$ composite function property depends only on the image X of φ , which is a closed subanalytic set [1] (cf. Corollary 1.5 below). Bierstone and Milman have proved, more generally, that a closed “Nash subanalytic” set X has the $\mathcal{C}^\infty$ composite function property [1] (cf. [19,23,26]); the class of Nash subanalytic sets

1980 *Mathematics Subject Classification* (1985 *Revision*). Primary 32B20, 58C27; Secondary 32K15, 58C25.

The first two authors' research was partially supported by NSERC operating grants OGP 0009070 and OGP 0008949.

includes all semianalytic sets. The $\mathcal{C}^\infty$ composite function property is equivalent to several other natural geometric and algebraic conditions on a closed subanalytic set [6]; in particular, to a formal semicoherence property (a stratified real version of the coherence theory of Oka and Cartan). Pawłucki has constructed an example of a closed subanalytic set which is not semicoherent [20]. Thus the $\mathcal{C}^\infty$ composite function property does not hold in general, but distinguishes an important class of subanalytic sets.

In this article, we introduce a new point of view towards Glaeser's theorem, with respect to which we can formulate a " $\mathcal{C}^k$ composite function property" that is satisfied by all semiproper real analytic mappings (Theorems 1.2 and 1.3 below). As a consequence, we see that a closed subanalytic set X satisfies the $\mathcal{C}^\infty$ composite function property if and only if the ring $\mathcal{C}^\infty(X)$ of $\mathcal{C}^\infty$ functions on X is the intersection of all finite differentiability classes (Corollary 1.5).

Let $k \in \mathbb{N} \cup \{\infty\}$, where $\mathbb{N}$ denotes the nonnegative integers. Suppose that A is a locally closed subset of $\mathbb{R}^n$ and that $B \subset A$ is closed (in the relative topology of A). We let $\mathcal{C}^k(A; B)$ denote the Fréchet algebra of restrictions to A of $\mathcal{C}^k$ real-valued functions that are defined on neighbourhoods of A and are k -flat on B . (" k -flat on B " means "vanishing on B together with all partial derivatives of orders $\leq k$ ".) $\mathcal{C}^k(A)$ means $\mathcal{C}^k(A; \emptyset)$.

Recall that a continuous mapping $\varphi: M \rightarrow N$ is called *semiproper* if, for each compact subset K of N , there is a compact subset L of M such that $\varphi(L) = K \cap \varphi(M)$. Let $\varphi: \Omega \rightarrow \mathbb{R}^n$ denote a semiproper real analytic mapping defined on an open subset Ω of $\mathbb{R}^m$. Then $X = \varphi(\Omega)$ is a closed subanalytic subset of $\mathbb{R}^n$. (For the definitions and basic properties of subanalytic and Nash subanalytic sets, see [1,5,7,8,9,14,16].) Let Z denote a closed subanalytic subset of X . The mapping φ induces injective homomorphisms $\varphi^*: \mathcal{C}^k(X; Z) \rightarrow \mathcal{C}^k(\Omega; \varphi^{-1}(Z))$ given by composition $\varphi^*(g) = g \circ \varphi$.

Definition 1.1. Let $(\varphi^*\mathcal{C}^k(X))^\wedge$ denote the subalgebra of all functions $f \in \mathcal{C}^k(\Omega)$ such that f is “formally a composite with φ ”; i.e., for each $a \in X$, there is $g \in \mathcal{C}^k(X)$ such that $f - \varphi^*(g)$ is k -flat on $\varphi^{-1}(a)$. Set $(\varphi^*\mathcal{C}^k(X; Z))^\wedge = (\varphi^*\mathcal{C}^k(X))^\wedge \cap \mathcal{C}^k(\Omega; \varphi^{-1}(Z))$.

Theorem 1.2. Assume that X is compact. Then, for each $k \in \mathbb{N}$, there is an integer $\ell_k \geq k$ such that $(\varphi^*\mathcal{C}^\ell(X; Z))^\wedge \subset \varphi^*\mathcal{C}^k(X; Z)$ for all $\ell \geq \ell_k$.

Glaeser’s theorem follows from Theorem 1.2: Assume that φ is generically a submersion, so that the interior $\text{int } X$ of X is dense in X . Suppose that $f \in (\varphi^*\mathcal{C}^\infty(X))^\wedge$. Let $g \in \mathcal{C}^0(X)$ denote the unique function such that $f = g \circ \varphi$. Then g is $\mathcal{C}^\infty$ in $\text{int } X$ and, by Theorem 1.2, g is the restriction to X of a $\mathcal{C}^k$ function, for all $k \in \mathbb{N}$. Thus all partial derivatives of $g|_{\text{int } X}$ extend continuously to X , and define a $\mathcal{C}^\infty$ Whitney field on X (cf. §3 below). By Whitney’s extension theorem, g is the restriction to X of a $\mathcal{C}^\infty$ function.

In general, put $\mathcal{C}^{(\infty)}(X; Z) = \bigcap_{k \in \mathbb{N}} \mathcal{C}^k(X; Z)$ (and $\mathcal{C}^{(\infty)}(X) = \mathcal{C}^{(\infty)}(X; \emptyset)$). The formula $\varphi^*(g) = g \circ \varphi$ defines an injective homomorphism $\varphi^*: \mathcal{C}^{(\infty)}(X; Z) \rightarrow \mathcal{C}^\infty(\Omega; \varphi^{-1}(Z))$. It is easy to see that $\bigcap_{\ell \in \mathbb{N}} (\varphi^*\mathcal{C}^\ell(X; Z))^\wedge = (\varphi^*\mathcal{C}^\infty(X; Z))^\wedge$, so that Theorem 1.2 implies the following:

Theorem 1.3. $(\varphi^*\mathcal{C}^\infty(X; Z))^\wedge = \varphi^*\mathcal{C}^{(\infty)}(X; Z)$.

There is a continuous injection $\mathcal{C}^\infty(X; Z) \hookrightarrow \mathcal{C}^{(\infty)}(X; Z)$, where the latter has a topology as the inverse limit of the $\mathcal{C}^k(X, Z)$, $k \in \mathbb{N}$; it is easy to see that $\mathcal{C}^\infty(X; Z)$ is dense in $\mathcal{C}^{(\infty)}(X; Z)$ (cf. [18, I.4.3]). Moreover, $(\varphi^*\mathcal{C}^\infty(X; Z))^\wedge$ is closed in $\mathcal{C}^\infty(\Omega; \varphi^{-1}(Z))$, so that Theorem 1.3 has the following corollaries:

Corollary 1.4. (cf. Tougeron [24]). $(\varphi^*\mathcal{C}^\infty(X; Z))^\wedge = \overline{\varphi^*\mathcal{C}^\infty(X; Z)}$.

Corollary 1.5. The subalgebra $\varphi^*\mathcal{C}^\infty(X; Z)$ is closed in $\mathcal{C}^\infty(\Omega; \varphi^{-1}(Z))$ if and only if $\mathcal{C}^{(\infty)}(X; Z) = \mathcal{C}^\infty(X; Z)$. In particular, closedness of $\varphi^*\mathcal{C}^\infty(X)$ in $\mathcal{C}^\infty(\Omega)$

depends only on the image X of φ (cf. [1,6]).

Theorem 1.6. *Let X be a closed subanalytic subset of $\mathbb{R}^n$, and let $Z \subset X$ denote the points which admit no neighbourhood in which X is Nash. (By [21], Z is a closed subanalytic set.) Then $\mathcal{C}^{(\infty)}(X; Z) = \mathcal{C}^\infty(X; Z)$.*

Theorem 1.6 can be obtained by combining Corollary 1.5 with [6, Theorem 3] and [22, Theorem 2]. In particular, if X is a closed Nash subanalytic subset of $\mathbb{R}^n$, then $\mathcal{C}^{(\infty)}(X) = \mathcal{C}^\infty(X)$. The same conclusion holds if $\dim X \leq 2$ or if X has pure codimension 1 (by [22]). It follows that, for any closed subanalytic subset X of $\mathbb{R}^4$, $\mathcal{C}^{(\infty)}(X) = \mathcal{C}^\infty(X)$. The example of [20] is a closed 3-dimensional subanalytic subset X of $\mathbb{R}^5$ for which $\mathcal{C}^{(\infty)}(X) \neq \mathcal{C}^\infty(X)$.

Remark 1.7. Theorems 1.2, 1.3 and their corollaries have more general module versions; i.e., versions that apply not only to the solution of an equation of the form $f(x) = g(\varphi(x))$ as above (where g is the unknown), but also more generally to systems of equations of the form $f(x) = A(x) \cdot g(\varphi(x)) + B(x) \cdot h(x)$, where $A(x)$, $B(x)$ are given matrices of real analytic functions, $f(x)$ is a given (vector-valued) $\mathcal{C}^k$ function, and $g(y)$, $h(x)$ are the unknowns (cf. [3,4]).

Remark 1.8. The results above can be stated, more generally, for a semiproper mapping $\varphi: M \rightarrow N$ of real analytic manifolds: The reduction to $N = \mathbb{R}^n$ is immediate. On the other hand, if $X = \varphi(M)$ is compact, then there are finitely many coordinate charts Ω_j in M such that $X = \cup \varphi(\Omega_j)$ and φ is semiproper when regarded as a mapping from the disjoint union Ω of the Ω_j . Therefore, we can reduce the statement of Theorem 1.2 for $\varphi: M \rightarrow N$ to the theorem as stated above.

2. DIVISION IN RINGS OF FORMAL POWER SERIES

We will prove Theorem 1.2 by the method used in [3,4] based on the division theorem of Grauert and Hironaka.

We totally order $\mathbb{N}^n$ using the lexicographic ordering of $(n+1)$ -tuples $(|\alpha|, \alpha_1, \dots, \alpha_n)$, where $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{N}^n$ and $|\alpha| = \alpha_1 + \dots + \alpha_n$. For every formal power series $F = \sum_{\alpha} f_{\alpha} y^{\alpha} \in \mathbb{R}[[y]] = \mathbb{R}[[y_1, \dots, y_n]]$, we define the *support* $\text{supp } F = \{\alpha \in \mathbb{N}^n : f_{\alpha} \neq 0\}$, the *initial exponent* $\exp F = \min \text{supp } F$ ($\exp F = \infty$ if $F = 0$), and the *initial monomial* $\text{mon } F = f_{\alpha} y^{\alpha}$, where $\alpha = \exp F$ (when $F \neq 0$).

Theorem 2.1. (cf. [11,15]). Let $F_1, \dots, F_s \in \mathbb{R}[[y]] \setminus \{0\}$, and let $\alpha^i = \exp F_i$ ($i = 1, \dots, s$). Put $\Delta_0 = \emptyset$, $\Delta_i = (\alpha^i + \mathbb{N}^n) \setminus (\bigcup_{j < i} \Delta_j)$ for $i = 1, \dots, s$, and $\Delta = \mathbb{N}^n \setminus (\bigcup_{i=0}^s \Delta_i)$. Let $G \in \mathbb{R}[[y]]$. Then there are unique $Q_i, R \in \mathbb{R}[[y]]$ ($i = 1, \dots, s$) such that $\alpha^i + \text{supp } Q_i \subset \Delta_i$ ($i = 1, \dots, s$), $\text{supp } R \subset \Delta$, and

$$G = \sum_{i=1}^s Q_i F_i + R.$$

Moreover, $\exp R \geq \exp G$ and, for each i , $\alpha^i + \exp Q_i \geq \exp G$.

See [4, p. 207] for the proof.

Let I be an ideal in $\mathbb{R}[[y]]$. The *diagram of initial exponents* of I is defined as $\mathfrak{N}(I) = \{\exp F : F \in I \setminus \{0\}\}$. Clearly, $\mathfrak{N}(I) + \mathbb{N}^n = \mathfrak{N}(I)$. This property implies that there is a smallest finite set $\mathfrak{V}(I) \subset \mathfrak{N}(I)$ such that $\mathfrak{V}(I) + \mathbb{N}^n = \mathfrak{N}(I)$. The elements of $\mathfrak{V}(I)$ are called the *vertices* of $\mathfrak{N}(I)$.

Let $\mathfrak{V}(I) = \{\alpha^1, \dots, \alpha^s\}$, and let $\{\Delta_i, \Delta\}$ denote the decomposition of $\mathbb{N}^n$ determined by the α^i , as in Theorem 2.1.

Proposition 2.2. [4, Corollary 6.8]. Choose $F_1, \dots, F_s \in I$ such that $\exp F_i = \alpha^i$ ($i = 1, \dots, s$). Then $F_1, \dots, F_s$ generate I . Moreover there is a unique

system $\{G_1, \dots, G_s\}$ of generators of I such that, for each i , $\text{supp}(G_i - y^{\alpha^i}) \subset \Delta$; in particular, $\text{mon} G_i = y^{\alpha^i}$.

We call $\{G_1, \dots, G_s\}$ the *standard basis* of I . Using Theorem 2.1, we get:

Proposition 2.3. (cf. [4, Corollary 6.9]). Let (y) denote the maximal ideal of $\mathbb{R}[[y]]$. Let $p \in \mathbb{N}$. Then the set $\{y^\beta : \beta \in \Delta, |\beta| \leq p\}$ is a basis of an $\mathbb{R}$ -linear complement of $I + (y)^{p+1}$ in $\mathbb{R}[[y]]$. In particular, if $(y)^{p+1} \subset I$, then $\{y^\beta : \beta \in \Delta\}$ is a basis of an $\mathbb{R}$ -linear complement of I in $\mathbb{R}[[y]]$.

Let $\mathcal{D}(n) = \{\mathfrak{N} \subset \mathbb{N}^n : \mathfrak{N} + \mathbb{N}^n = \mathbb{N}^n\}$. We totally order $\mathcal{D}(n)$ as follows: To each $\mathfrak{N} \in \mathcal{D}(n)$, associate the sequence $v(\mathfrak{N})$ obtained by listing the vertices of $\mathfrak{N}$ in ascending order and completing the list to an infinite sequence by using ∞ for all the remaining terms. If $\mathfrak{N}_1, \mathfrak{N}_2 \in \mathcal{D}(n)$, we say that $\mathfrak{N}_1 < \mathfrak{N}_2$ provided that $v(\mathfrak{N}_1) < v(\mathfrak{N}_2)$ with respect to the lexicographic ordering on the set of all such sequences.

3. WHITNEY FIELDS

Let $p \in \mathbb{N} \setminus \{0\}$ and let A be a locally closed subset of $\mathbb{R}^n$. A $\mathcal{C}^p$ -Whitney field on A is a polynomial $F(a, y) = \sum_{|\alpha| \leq p} \frac{1}{\alpha!} F_\alpha(a) y^\alpha \in \mathcal{C}^0(A)[y] = \mathcal{C}^0(A)[y_1, \dots, y_n]$ which fulfills the following conditions:

$$\left| \left(\partial^{|\beta|} F / \partial y^\beta \right) (a, 0) - \left(\partial^{|\beta|} F / \partial y^\beta \right) (b, a - b) \right| / |a - b|^{p-|\beta|} \longrightarrow 0,$$

when $a \rightarrow c$, $b \rightarrow c$, $a \neq b$, for each $c \in A$ and $\beta \in \mathbb{N}^n$ such that $|\beta| \leq p$.

For any polynomial $F(a, y) = \sum_{|\alpha| \leq p} \frac{1}{\alpha!} F_\alpha(a) y^\alpha \in \mathcal{C}^0(A)[y]$, we set $F^k(a, y) = \sum_{|\alpha| \leq k} \frac{1}{\alpha!} F_\alpha(a) y^\alpha$, where $k \leq p$, $k \in \mathbb{N}$.

For any $\mathcal{C}^p$ -mapping $\varphi = (\varphi_1, \dots, \varphi_n) : D \rightarrow \mathbb{R}^n$ defined on an open subset D of $\mathbb{R}^m$, we denote by $T^p \varphi$ its *Taylor field*

$$T^p \varphi(b, x) = \sum_{|\alpha| \leq p} \frac{1}{\alpha!} D^\alpha \varphi(b) x^\alpha = \left(\sum_{|\alpha| \leq p} \frac{1}{\alpha!} D^\alpha \varphi_1(b) x^\alpha, \dots, \sum_{|\alpha| \leq p} \frac{1}{\alpha!} D^\alpha \varphi_n(b) x^\alpha \right).$$

Put $\tilde{T}^p\varphi(b, x) = T^p\varphi(b, x) - \varphi(b)$ (the field of Taylor polynomials without constant terms).

Lemma 3.1. *Let Λ be a $\mathcal{C}^1$ -submanifold of $\mathbb{R}^n$. Let $F(a, y) = \sum_{|\alpha| \leq p} \frac{1}{\alpha!} F_\alpha(a) y^\alpha \in \mathcal{C}^0(\Lambda)[y]$, where $F^{p-1}(a, y) \in \mathcal{C}^1(\Lambda)[y]$. Let $\varphi : D \rightarrow \mathbb{R}^n$ be a $\mathcal{C}^p$ -mapping as above, and let Γ be a $\mathcal{C}^1$ -submanifold of D such that $\varphi(\Gamma) \subset \Lambda$. Define a polynomial $G \in \mathcal{C}^0(\Gamma)[x] = \mathcal{C}^0(\Gamma)[x_1, \dots, x_m]$ of degree $\leq p$, by the formula*

$$G(b, x) = F(\varphi(b), \tilde{T}^p\varphi(b, x)) \bmod (x)^{p+1}.$$

Then $G^{p-1} \in \mathcal{C}^1(\Gamma)[x]$ and, for each $b \in \Gamma$ and $v \in T_b\Gamma$, we have

$$\begin{aligned} D_{b,v}G^{p-1}(b, x) - D_{x,v}G(b, x) \\ = D_{a,u}F^{p-1}(\varphi(b), \tilde{T}^p\varphi(b, x)) - D_{y,u}F(\varphi(b), \tilde{T}^p\varphi(b, x)) \bmod (x)^p, \end{aligned}$$

where $u = d_b\varphi(v)$, and $D_{b,v}$ and $D_{x,v}$ stand for the (directional) derivatives with respect to b and x , respectively, along the tangent vector v .

This is a straightforward calculation.

Proposition 3.2. (cf. [4, §10]). *Let Λ be a $\mathcal{C}^p$ -submanifold of $\mathbb{R}^n$ and let $F \in \mathcal{C}^0(\Lambda)[y]$, $\deg F \leq p$. Then F is a $\mathcal{C}^p$ -Whitney field on Λ if and only if $F^{p-1} \in \mathcal{C}^1(\Lambda)[y]$ and $D_{a,u}F^{p-1}(a, y) = D_{y,u}F(a, y)$, for every $a \in \Lambda$ and $u \in T_a\Lambda$.*

Proof. It is enough to consider the case $\Lambda = \mathbb{R}^k \times 0$; the proposition then follows easily. $\square$

Remark 3.3. [25, IV.2.5]. Let U denote an open subset of $\mathbb{R}^n$. Let $u, v \in U$, and suppose that σ is a rectifiable curve in U joining u, v . Let $g \in \mathcal{C}^k(U)$, $k \geq 1$. It follows from the Mean Value Theorem (using an approximation of σ by piecewise-linear curves) that

$$|g(u) - g(v)| \leq \sqrt{n}|\sigma| \sup_{\substack{a \in \sigma \\ 1 \leq j \leq n}} \left| \frac{\partial g}{\partial y_j}(a) \right|,$$

where $|\sigma|$ denotes the length of σ . If g is $(k-1)$ -flat at v , then (by iterating the inequality above) we get

$$|g(u)| \leq n^{k/2} |\sigma|^k \sup_{\substack{a \in \sigma \\ |\alpha|=k}} \left| \frac{\partial^{|\alpha|} g}{\partial y^\alpha}(a) \right|.$$

Now suppose that F is a $\mathcal{C}^\ell$ Whitney field on σ . Let $\alpha \in \mathbb{N}$, $|\alpha| < \ell$, and apply the preceding inequality with $k = \ell - |\alpha|$ and $g(y) = \frac{\partial^{|\alpha|}}{\partial y^\alpha}(f(y) - F(v, y - v))$, where $f \in \mathcal{C}^\ell(U)$ is a Whitney extension of F . We obtain

$$\left| \frac{\partial^{|\alpha|} F}{\partial y^\alpha}(u, 0) - \frac{\partial^{|\alpha|} F}{\partial y^\alpha}(v, u - v) \right| \leq n^{\frac{\ell-|\alpha|}{2}} |\sigma|^{\ell-|\alpha|} \sup_{\substack{a \in \sigma \\ |\beta|=\ell}} |F_\beta(a) - F_\beta(v)|.$$

(This inequality holds trivially if $|\alpha| = \ell$).

Let $r \in \mathbb{N} \setminus \{0\}$. A compact subset A of $\mathbb{R}^n$ is called *r-regular* if there is a constant $C > 0$ such that any two points $u, v \in A$ can be joined by a rectifiable curve σ in A of length $|\sigma| \leq C|u - v|^{1/r}$. The following is a version of l'Hôpital's rule or Hestenes's lemma for *r-regular* sets (generalizing [27]).

Proposition 3.4. *Let $A \supset B$ be compact subsets of $\mathbb{R}^n$, where A is *r-regular*. Let $k \in \mathbb{N}$ and let $F(a, y) = \sum_{|\alpha| \leq kr} \frac{1}{\alpha!} F_\alpha(a) y^\alpha \in \mathcal{C}^0(A)[y]$, $y = (y_1, \dots, y_n)$. Assume that F restricts to $\mathcal{C}^{kr}$ Whitney fields on B and on $A \setminus B$. Then $F^k(a, y) = \sum_{|\alpha| \leq k} \frac{1}{\alpha!} F_\alpha(a) y^\alpha$ is a $\mathcal{C}^k$ Whitney field on A .*

Proof. For all $u, v \in A$ and each $\alpha \in \mathbb{N}^n$, $|\alpha| \leq k$,

$$\begin{aligned} & \frac{\partial^{|\alpha|} F^k}{\partial y^\alpha}(u, 0) - \frac{\partial^{|\alpha|} F^k}{\partial y^\alpha}(v, u - v) \\ (3.5) \quad &= \frac{\partial^{|\alpha|} F}{\partial y^\alpha}(u, 0) - \frac{\partial^{|\alpha|} F}{\partial y^\alpha}(v, u - v) + \sum_{k-|\alpha| < |\beta| \leq kr-|\alpha|} \frac{1}{\beta!} F_{\alpha+\beta}(v) (u - v)^\beta. \end{aligned}$$

We can assume that F is kr -flat on B (by Whitney's extension theorem). Let $u, v \in A$, and let σ be a rectifiable curve in A of length $|\sigma| \leq C|u - v|^{1/r}$ joining u, v . Write

$$\mu_F(\sigma) = \sup_{\substack{a, b \in \sigma \\ |\beta|=kr}} |F_\beta(a) - F_\beta(b)|,$$

so that $\mu_F(\sigma) \rightarrow 0$ as $|u - v| \rightarrow 0$.

Case 1. $\sigma \cap B = \emptyset$. By Remark 3.3, for all $|\alpha| \leq kr$,

$$\begin{aligned} \left| \frac{\partial^{|\alpha|} F}{\partial y^\alpha}(u, 0) - \frac{\partial^{|\alpha|} F}{\partial y^\alpha}(v, u - v) \right| &\leq n^{\frac{kr - |\alpha|}{2}} |\sigma|^{kr - |\alpha|} \mu_F(\sigma) \\ &\leq c_1 |u - v|^{k - |\alpha|} \mu_F(\sigma), \end{aligned}$$

where c_1 is independent of u, v . By (3.5), if $|\alpha| \leq k$, then

$$\left| \frac{\partial^{|\alpha|} F^k}{\partial y^\alpha}(u, 0) - \frac{\partial^{|\alpha|} F^k}{\partial y^\alpha}(v, u - v) \right| \leq c_2 |u - v|^{k - |\alpha|} (\mu_F(\sigma) + |u - v|).$$

Case 2. $\sigma \cap B \neq \emptyset$. If $w \in \sigma$, let $\sigma_{u,w}$ denote the shortest part of σ joining u, w . Then there exists $w \in \sigma$ such that $\sigma_{u,w} \cap B = \{w\}$. Using the Mean Value Theorem (as in Remark 3.3), we see that if $|\alpha| \leq kr$, then

$$\begin{aligned} (3.6) \quad |F_\alpha(u)| &\leq n^{\frac{kr - |\alpha|}{2}} |\sigma_{u,w}|^{kr - |\alpha|} \sup_{\substack{a \in \sigma_{u,w} \\ |\beta| = kr}} |F_\beta(a)| \\ &\leq n^{\frac{kr - |\alpha|}{2}} |\sigma|^{kr - |\alpha|} \mu_F(\sigma) \\ &\leq c_1 |u - v|^{k - |\alpha|} \mu_F(\sigma), \end{aligned}$$

and (applying (3.6) with v in place of u)

$$\left| \sum_{|\beta| \leq kr - |\alpha|} \frac{F_{\alpha+\beta}(v)}{\beta!} (u - v)^\beta \right| \leq c_3 |u - v|^{k - |\alpha|} \mu_F(\sigma).$$

Thus

$$\left| \frac{\partial^{|\alpha|} F}{\partial y^\alpha}(u, 0) - \frac{\partial^{|\alpha|} F}{\partial y^\alpha}(v, u - v) \right| \leq (c_1 + c_3) |u - v|^{k - |\alpha|} \mu_F(\sigma),$$

and the required estimate again follows from (3.5). $\square$

4. STRATIFICATION OF A SUBANALYTIC MAPPING

We will use a theorem of Hardt on stratification of mappings [12,13] in the version of Łojasiewicz [17].

A *subanalytic leaf* in $\mathbb{R}^m$ means a connected subanalytic subset of $\mathbb{R}^m$ which is an analytic submanifold. Let E be a subanalytic subset of $\mathbb{R}^m$. A *subanalytic*

stratification of E (in $\mathbb{R}^m$) is a partition $\mathcal{S}$ of E into subanalytic leaves in $\mathbb{R}^m$, called *strata*, locally finite in $\mathbb{R}^m$, such that, for each $S \in \mathcal{S}$, the boundary $(\overline{S} \setminus S) \cap E$ is a union of strata of dimension $< \dim S$. A mapping $\varphi: E \rightarrow \mathbb{R}^n$ is called *subanalytic* if its graph is subanalytic in $\mathbb{R}^m \times \mathbb{R}^n$. A partition $\mathcal{P}$ of E is called *compatible* with a family $\mathcal{F}$ of subsets of E if, for each $P \in \mathcal{P}$ and $F \in \mathcal{F}$, either $P \subset F$ or $P \subset E \setminus F$.

Theorem 4.1. (cf. [17]). *Let $\varphi: E \rightarrow \mathbb{R}^n$ be a continuous subanalytic mapping defined on a compact subanalytic subset E of $\mathbb{R}^m$. Put $X = \varphi(E)$. Let $\mathcal{F}$ and $\mathcal{G}$ be finite families of subsets of E and X which are subanalytic in $\mathbb{R}^m$ and $\mathbb{R}^n$, respectively. Then there exist finite subanalytic stratifications $\mathcal{S}$ and $\mathcal{T}$ of E and X , respectively, such that:*

(1) *For each $S \in \mathcal{S}$, $\varphi(S) \in \mathcal{T}$ and there is a commutative diagram*

$$\begin{array}{ccc} S & \xrightarrow{h} & T \times P \\ \varphi|_S \searrow & & \swarrow \pi \\ & T & \end{array}$$

where $T = \varphi(S)$, P is a bounded subanalytic leaf in $\mathbb{R}^s$ for some s , h is an analytic subanalytic isomorphism and π denotes the natural projection.

(2) *$\mathcal{S}$ is compatible with $\mathcal{F}$ and $\mathcal{T}$ is compatible with $\mathcal{G}$.*

A pair $(\mathcal{S}, \mathcal{T})$ of subanalytic stratifications as in (1) will be called a *stratification* of φ . If we weaken this definition by allowing $\mathcal{S}$ to be any finite partition into subanalytic leaves (but still requiring $\mathcal{T}$ to be a stratification), then the pair $(\mathcal{S}, \mathcal{T})$ will be called a *semistratification* of φ .

Remark 4.2. Clearly, Theorem 4.1 is true for a subanalytic mapping $\varphi: E \rightarrow \mathbb{R}^n$ which is defined on a bounded subanalytic subset E of $\mathbb{R}^m$ and extends continuously to $\overline{E}$.

Remark 4.3. If $(\mathcal{S}, \mathcal{T})$ is a semistratification of φ and Y is a subanalytic subset of X such that $\mathcal{T}' = \{T \cap Y : T \in \mathcal{T}\}$ is a stratification of Y , then the pair

$(\mathcal{S}', \mathcal{T}')$, where $\mathcal{S}' = \{S \cap \varphi^{-1}(Y) : S \in \mathcal{S}\}$, is a semistratification of $\varphi|_{\varphi^{-1}(Y)} : \varphi^{-1}(Y) \rightarrow Y$.

Remark 4.4. If $(\mathcal{S}, \mathcal{T})$ is a semistratification of φ , $S \in \mathcal{S}$, $T \in \mathcal{T}$ and $\varphi(S) = T$, then there exists a subanalytic leaf $\Gamma \subset S$ such that $\varphi|_{\Gamma} : \Gamma \rightarrow T$ is an analytic isomorphism.

Let $\varphi : E \rightarrow \mathbb{R}^n$ be a bounded subanalytic mapping defined on a bounded subanalytic subset E of $\mathbb{R}^m$. For each $q \in \mathbb{N}$, $q \geq 1$, we define the q -fold fibre product of E with respect to φ ,

$$E_\varphi^q = \{\underline{b} = (b_1, \dots, b_q) \in E^q : \varphi(b_1) = \dots = \varphi(b_q)\} ;$$

E_φ^q is a subanalytic subset of $(\mathbb{R}^m)^q$. There is a natural mapping $\Phi : E_\varphi^q \rightarrow \mathbb{R}^n$ defined by $\Phi(b_1, \dots, b_q) = \varphi(b_1)$. Suppose that $(\mathcal{S}, \mathcal{T})$ is a semistratification of φ . Let $\mathcal{S}^{(q)}$ denote the family of all non-empty sets of the form $(S_1 \times \dots \times S_q) \cap E_\varphi^q$, where $S_1, \dots, S_q \in \mathcal{S}$. It is easy to see (using Remark 4.3) that $(\mathcal{S}^{(q)}, \mathcal{T})$ is a semistratification of Φ .

5. IDEALS OF RELATIONS

Let E be a bounded open subanalytic subset of $\mathbb{R}^m$, and let $\varphi = (\varphi_1, \dots, \varphi_n) : E \rightarrow \mathbb{R}^n$ be a mapping which extends to be analytic in a neighbourhood of $\overline{E}$. Let $p \in \mathbb{N}$, $p \geq 1$. Put $X = \varphi(E)$ and $q = \binom{n+p}{p}$.

For each $a \in X$, we define the *ideal of relations of order p among $\varphi_1, \dots, \varphi_n$ over a* as

$$\mathcal{R}_p(a) = \{W \in \mathbb{R}[[y]] : W(\tilde{T}^p \varphi(b, x)) = 0 \text{ mod } (x)^{p+1}, \text{ for each } b \in \varphi^{-1}(a)\}$$

Clearly, $(y)^{p+1} \subset \mathcal{R}_p(a)$. Put

$$L_\beta^\alpha(b) = D_x^\beta((\tilde{T}^p \varphi(b, x))^\alpha)(0) ,$$

where $\alpha \in \mathbb{N}^n$, $\beta \in \mathbb{N}^m$, $|\alpha| \leq p$, $|\beta| \leq p$, $b \in E$. (D_x^β denotes the partial derivative $\partial^{|\beta|}/\partial x^\beta$ with respect to the variables x .) Then for every formal power series $W = \sum_\alpha W_\alpha y^\alpha \in \mathbb{R}[[y]]$, $W \in \mathcal{R}_p(a)$ if and only if $\{W_\alpha\}_{|\alpha| \leq p}$ satisfies the following system of linear equations:

$$(5.1) \quad \sum_{|\alpha| \leq p} W_\alpha \cdot L_\beta^\alpha(b) = 0 \quad (|\beta| \leq p, b \in \varphi^{-1}(a)).$$

Let $(\mathcal{S}, \mathcal{T})$ be a semistratification of φ . This induces a semistratification $(\mathcal{S}^{(q)}, \mathcal{T})$ of the mapping $\Phi: E_\varphi^q \rightarrow X$. Let $l \in \mathbb{N}$, $1 \leq l \leq p$. Denote by $\pi_l: \mathbb{R}[[y]] \rightarrow \mathbb{R}[y]$ the projection $\pi_l(\sum_\alpha W_\alpha y^\alpha) = \sum_{|\alpha| \leq l} W_\alpha y^\alpha$. It is clear that $\pi_l(\mathcal{R}_r(a)) \subset \pi_l(\mathcal{R}_p(a)) \subset \mathcal{R}_l(a)$, when $l \leq p \leq r$ and $a \in X$.

For each $\underline{b} = (b_1, \dots, b_q) \in E_\varphi^q$, let $\rho^0(\underline{b})$ denote the rank of the matrix of the system

$$(5.2) \quad \sum_{|\alpha| \leq p} W_\alpha L_\beta^\alpha(b_\nu) = 0 \quad (|\beta| \leq p, \nu = 1, \dots, q),$$

and $\rho^1(\underline{b})$ the rank of the matrix of the system

$$(5.3) \quad \sum_{l < |\alpha| \leq p} W_\alpha L_\beta^\alpha(b_\nu) = 0 \quad (|\beta| \leq p, \nu = 1, \dots, q).$$

(Of course, $\rho^0(\underline{b})$ depends on p , and $\rho^1(\underline{b})$ on p and l .) For each $T \in \mathcal{T}$, we put

$$\sigma_T^0 = \max\{\rho^0(\underline{b}) : \underline{b} \in \Phi^{-1}(T)\},$$

$$\sigma_T^1 = \max\{\rho^1(\underline{b}) : \underline{b} \in \Phi^{-1}(T)\}.$$

Observe that if $a \in T$ is a point such that

$$\max\{\rho^0(\underline{b}) : \underline{b} \in \Phi^{-1}(a)\} = \sigma_T^0,$$

$$\max\{\rho^1(\underline{b}) : \underline{b} \in \Phi^{-1}(a)\} = \sigma_T^1,$$

then $\dim \pi_p(\mathcal{R}_p(a)) = q - \sigma_T^0$ and $\dim \pi_l(\mathcal{R}_p(a)) = q - \sigma_T^0 - (q - \binom{l+n}{n}) - \sigma_T^1 = \binom{l+n}{n} + \sigma_T^1 - \sigma_T^0$.

For each $T \in \mathcal{T}$, let us take leaves $S_T^0, S_T^1 \in \mathcal{S}^{(q)}$ such that $\Phi(S_T^0) = \Phi(S_T^1) = T$ and

$$\max\{\rho^0(\underline{b}) : \underline{b} \in S_T^0\} = \sigma_T^0,$$

$$\max\{\rho^1(\underline{b}) : \underline{b} \in S_T^1\} = \sigma_T^1.$$

Then there is a closed nowhere dense subset $Z_T^{p_l}$ of T , subanalytic in $\mathbb{R}^n$, such that, for each $a \in T \setminus Z_T^{p_l}$,

$$\max\{\rho^0(\underline{b}) : \underline{b} \in S_T^0 \cap \Phi^{-1}(a)\} = \sigma_T^0,$$

$$\max\{\rho^1(\underline{b}) : \underline{b} \in S_T^1 \cap \Phi^{-1}(a)\} = \sigma_T^1.$$

Put $\omega_T^{p_l} = \binom{l+n}{n} + \sigma_T^1 - \sigma_T^0$. Then $\omega_T^{p_l} = \dim \pi_l(\mathcal{R}_p(a))$, for $a \in T \setminus Z_T^{p_l}$. We have $\omega_T^{r_l} \leq \omega_T^{p_l}$, when $l \leq p \leq r$. Therefore, for every l , there exists $p_l \geq l$ such that $\omega_T^{p_l}$ is constant for $p \geq p_l$. Since $\mathcal{T}$ is finite, we can make p_l independent of T . This gives the following:

Lemma 5.4. *For every positive $l \in \mathbb{N}$, there exists $p_l \in \mathbb{N}$, $p_l \geq l$, such that if $p > p_l$ and $T \in \mathcal{T}$, then $\pi_l(\mathcal{R}_p(a)) = \pi_l(\mathcal{R}_{p-1}(a))$, for each $a \in T \setminus (Z_T^{p_l} \cup Z_T^{(p-1)l})$.*

Now let us fix any $p > p_l$. Take $T \in \mathcal{T}$. The diagram of initial exponents $\mathfrak{N}(\mathcal{R}_p(a))$, $a \in T \setminus Z_T^{p_l}$, takes only finitely many values; choose $a \in T \setminus Z_T^{p_l}$ with the minimum value (cf. §2). Set $\Delta(a) = \mathbb{N}^n \setminus \mathcal{N}(\mathcal{R}_p(a))$. There exists $\underline{b} \in S_T^0 \cap \Phi^{-1}(a)$ such that $\rho^0(\underline{b}) = \sigma_T^0$. By Proposition 2.3, for any such $\underline{b}$, we have $\#\Delta(a) = \rho^0(\underline{b}) = \sigma_T^0 = \text{rank}\{L_{\beta}^{\alpha}(b_{\nu})\}$ (where $\alpha \in \Delta(a)$, $|\beta| \leq p$, $\nu = 1, \dots, q$). There exists a nonzero determinant

$$M_T(\underline{b}) = \det(\{L_{\beta_k}^{\alpha}(b_{\nu_k})\}) \quad (\alpha \in \Delta(a), k = 1, \dots, \sigma_T^0).$$

Put $\Theta_T^{p_l} = \{\underline{c} \in S_T^0 : M_T(\underline{c}) = 0\}$ and $\Sigma_T^{p_l} = T \setminus \Phi(S_T^0 \setminus \Theta_T^{p_l})$. Clearly, $\Theta_T^{p_l}$ and $\Sigma_T^{p_l}$ are nowhere dense in S_T^0 and in T , and are subanalytic in $(\mathbb{R}^m)^q$ and in $\mathbb{R}^n$, respectively. Since $M_T(\underline{b}) \neq 0$, it follows from Cramer's rule that the system

(5.2), and therefore the system (5.1), is equivalent to the system

$$(5.5) \quad W_\gamma - \sum_{\alpha \notin \Delta(a)} W_\alpha \cdot N_{T,\gamma}^\alpha(\underline{b})/M_T(\underline{b}) = 0 \quad (\gamma \in \Delta(a)),$$

where $N_{T,\gamma}^\alpha(\underline{b})$ are appropriate minors of the matrix of the system (5.2). Let $\{\alpha^1, \dots, \alpha^s\} = \mathfrak{V}(\mathcal{R}_p(a))$, where $\alpha^1 < \dots < \alpha^s$, and let $r = \max\{i : |\alpha^i| \leq p\}$. Since $(y)^{p+1} \subset \mathcal{R}_p(a)$, $|\alpha^i| = p+1$ for $i > r$. Observe that the standard basis of $\mathcal{R}_p(a)$ consists of y^{α^i} ($i = r+1, \dots, s$) and

$$G_i(y) = y^{\alpha^i} + \sum_{\gamma \in \Delta(a)} \left(N_{T,\gamma}^{\alpha^i}(\underline{b})/M_T(\underline{b}) \right) y^\gamma \quad (i = 1, \dots, r).$$

Moreover, we have

$$y^{\alpha^i} + \sum_{\gamma \in \Delta(a)} \left(N_{T,\gamma}^{\alpha^i}(\underline{b}')/M_T(\underline{b}') \right) y^\gamma \in \mathcal{R}_p(\Phi(\underline{b}')),$$

for any $\underline{b}' \in S_T^0 \setminus \Theta_T^{pl}$ such that $\Phi(\underline{b}') \notin Z_T^{pl}$, and $i = 1, \dots, r$. Suppose that $a' \in T \setminus (Z_T^{pl} \cup \Sigma_T^{pl})$. Since $\mathfrak{N}(\mathcal{R}_p(a)) \leq \mathfrak{N}(\mathcal{R}_p(a'))$, it follows that $\alpha^i = \exp G_i \in \mathfrak{N}(\mathcal{R}_p(a'))$ for each i , so that $\mathfrak{N}(\mathcal{R}_p(a)) \subset \mathfrak{N}(\mathcal{R}_p(a'))$ and hence they are equal. In other words, $\Delta(a')$ is independent of $a' \in T \setminus (Z_T^{pl} \cup \Sigma_T^{pl})$. Let us denote it by Δ_T .

Now taking an appropriate semistratification of Φ and using Remark 4.4, we can find a finite subanalytic stratification $\{\Lambda_j\}$ ($j \in J$) of X , compatible with the sets T , Z_T^{pl} , $Z_T^{(p-1)l}$, Σ_T^{pl} , and a family $\{\Gamma_j\}$ ($j \in J$) of subanalytic leaves in E_φ^q such that, for each $j \in J$, $\Phi|_{\Gamma_j} : \Gamma_j \rightarrow \Lambda_j$ is an analytic isomorphism, and $\Gamma_j \subset S_T^0 \setminus \Theta_T^{pl}$, whenever $\Lambda_j \subset T \setminus \Sigma_T^{pl}$.

Suppose that $j \in J$, $T \in \mathcal{T}$, $\Lambda_j \subset T$ and $\dim \Lambda_j = \dim T$. Then Λ_j is open in T and $\Lambda_j \subset T \setminus (Z_T^{pl} \cup \Sigma_T^{pl})$. We put $M_j = M_T$ and $\Delta_j = \Delta_T$ in this case.

6. PROOF OF THEOREM 1.2

Since φ is semiproper and $X = \varphi(\Omega)$ is compact, there exists a bounded open subanalytic subset $E \subset \mathbb{R}^m$ such that $\overline{E} \subset \Omega$, $\varphi(E) = X$ and $\varphi|_E : E \rightarrow \mathbb{R}^n$

is semiproper. Theorem 1.2 reduces to the following proposition (by induction on dimension).

Proposition 6.1. *Let A be a closed subset of X , subanalytic in $\mathbb{R}^n$, of dimension d . Let $k \in \mathbb{N}$. Then there exists a closed subset A' of A and an integer $t \geq k$ such that:*

- (1) A' is subanalytic of dimension $< d$.
- (2) If $f \in C^t(\Omega)$ is t -flat on $\varphi^{-1}(A' \cup Z)$ and, for each $a \in A$, there exists a polynomial $W_a \in \mathbb{R}[y]$ such that $T^t f(b, x) = W_a(\tilde{T}^t \varphi(b, x)) \bmod (x)^{t+1}$, for all $b \in \varphi^{-1}(a) \cap E$, then there exists $g \in C^k(\mathbb{R}^n; Z)$ such that $f - g \circ \varphi$ is k -flat on $\varphi^{-1}(A) \cap E$.

Proof. Let r denote a positive integer such that every connected component of $A \cup Z$ is r -regular [5, Th. 6.10]. Take any integer $l \geq kr$. Let $(\mathcal{S}, \mathcal{T})$ be a (semi) stratification of $\varphi|E$ such that $\mathcal{T}$ is compatible with A and Z . We will use the results and notation of §5. Fix an integer $p > p_l$ and take the stratification $\{\Lambda_j\}$ ($j \in J$) of X and the family $\{\Gamma_j\}$ ($j \in J$) of subanalytic leaves in E_φ^q as at the end of Section 5. Observe that the mapping $\Phi: E_\varphi^q \rightarrow X$ extends to $\Phi: \Omega^q \rightarrow X$, and $\overline{E_\varphi^q} \subset \Omega^q$. For each $j \in J$, $\Phi(\overline{\Gamma_j} \setminus \Gamma_j) = \overline{\Lambda_j} \setminus \Lambda_j$, because $\Phi|_{\Gamma_j}: \Gamma_j \rightarrow \Lambda_j$ is a homeomorphism.

Set $J_0 = \{j \in J: \Lambda_j \subset A \setminus Z, \dim \Lambda_j = d\}$ and $A' = \bigcup \{\Lambda_j: \Lambda_j \subset A, \dim \Lambda_j < d\}$. For each $j \in J_0$, Λ_j is contained in some $T \in \mathcal{T}$ of dimension d . Thus M_j is nonzero on Γ_j and we have a Łojasiewicz inequality

$$(6.2) \quad |M_j(\underline{b})| \geq C \cdot (\text{dist}(\underline{b}, \overline{\Gamma_j} \setminus \Gamma_j))^s,$$

for $\underline{b} \in \Gamma_j$, $j \in J_0$, where s is a positive integer.

Now take any integer $t \geq p + s$. Suppose that $f \in C^t(\Omega)$ is t -flat on $\varphi^{-1}(A' \cup Z)$ and that, for each $a \in A$, there exists a polynomial $W_a \in \mathbb{R}[y]$ such that $T^t f(b, x) = W_a(\tilde{T}^t \varphi(b, x)) \bmod (x)^{t+1}$ for each $b \in \varphi^{-1}(a) \cap E$.

Let $j \in J_0$. For each $a \in \Lambda_j$, let $V_j(a, y) \in \mathbb{R}[y]$ be the unique polynomial (of degree $\leq p$) such that $W_a(y) - V_j(a, y) \in \mathcal{R}_p(a)$ and $\text{supp}_y V_j(a, y) \subset \Delta_j$. Put

$$V_j(a, y) = \sum_{\alpha \in \Delta_j} V_j^\alpha(a) y^\alpha,$$

$$F_j(a, y) = \pi_l(V_j(a, y)) = \sum_{\substack{\alpha \in \Delta_j \\ |\alpha| \leq l}} V_j^\alpha(a) y^\alpha.$$

Lemma 6.3. $\lim V_j^\alpha(a) = 0$, when $a \rightarrow \bar{\Lambda}_j \setminus \Lambda_j$.

Proof. Since $T^p f(b, x) = V_j(a, \tilde{T}^p \varphi(b, x)) \bmod (x)^{p+1}$, for each $b \in \varphi^{-1}(a) \cap E$, we have

$$D^\beta f(b_\nu) = \sum_{\alpha \in \Delta_j} V_j^\alpha(\Phi(\underline{b})) L_\beta^\alpha(b_\nu)$$

where $\underline{b} = (b_1, \dots, b_q) \in \Gamma_j$, $\nu = 1, \dots, q$, $\beta \in \mathbb{N}^m$ and $|\beta| \leq p$. By Cramer's rule applied to this system,

$$(6.4) \quad V_j^\alpha(\Phi(\underline{b})) = H_j^\alpha(\underline{b}) / M_j(\underline{b})$$

for $\underline{b} \in \Gamma_j$, $\alpha \in \Delta_j$, where the H_j^α are $\mathcal{C}^s$ -functions on Ω^q , s -flat on $\bar{\Gamma}_j \setminus \Gamma_j \subset \Phi^{-1}(\bar{\Lambda}_j \setminus \Lambda_j) \subset \Phi^{-1}(A')$. The lemma follows from (6.2). $\square$

Lemma 6.5. F_j is a $\mathcal{C}^l$ -Whitney field on Λ_j .

Proof. Let $\Psi_\nu: \Omega^q \rightarrow \mathbb{R}^n$ and $h_\nu: \Omega^q \rightarrow \mathbb{R}$ ($\nu = 1, \dots, q$) denote the mappings $\Psi_\nu(\underline{b}) = \Psi_\nu(b_1, \dots, b_q) = \varphi(b_\nu)$ and $h_\nu(\underline{b}) = f(b_\nu)$. Then $\Psi_\nu|_{E_\varphi^q} = \Phi$ and, for each $\underline{b} \in \Omega^q$, the polynomial $T^p \Psi_\nu(\underline{b}, \underline{x})$ can be identified with $T^p \varphi(b_\nu, x)$. Similarly, $T^p h_\nu(\underline{b}, \underline{x})$ can be identified with $T^p f(b_\nu, x)$. Fix $a \in \Lambda_j$. Choose $\underline{b} \in \Phi^{-1}(a) \cap E_\varphi^q$ such that, for any $W \in \mathbb{R}[y]$, $W \in \mathcal{R}_{p-1}(a)$ if and only if $W(\tilde{T}^p \varphi(b_\nu, x)) = 0 \bmod (x)^p$, $\nu = 1, \dots, q$. Take $S \in \mathcal{S}^{(q)}$ such that $\underline{b} \in S$. Then Λ_j is open in $T = \Phi(S)$. We have

$$(6.6) \quad T^p h_\nu(\underline{c}, \underline{x}) = V_j\left(\Phi(\underline{c}), \tilde{T}^p \Psi_\nu(\underline{c}, \underline{x})\right) \bmod (\underline{x})^{p+1}$$

for all $\underline{c} \in S \cap \Phi^{-1}(\Lambda_j)$, $\nu = 1, \dots, q$. By (6.4), the V_j^α are $\mathcal{C}^1$ on Λ_j (in fact, $\mathcal{C}^s$). Let u be any vector tangent to Λ_j at a . Since $\Phi|_S: S \rightarrow T$ is a submersion, there is $v \in T_{\underline{b}}S$ such that $(d_{\underline{b}}(\Phi|_S))(v) = (d_{\underline{b}}\Psi_\nu)(v) = u$. By Lemma 3.1, Proposition 3.2 and (6.6),

$$D_{a,u}V_j^{p-1}\left(a, \tilde{T}^p\Psi_\nu(\underline{b}, \underline{x})\right) - D_{y,u}V_j\left(a, \tilde{T}^p\Psi_\nu(\underline{b}, \underline{x})\right) = 0 \bmod (\underline{x})^p;$$

hence

$$D_{a,u}V_j^{p-1}\left(a, \tilde{T}^p\varphi(b_\nu, x)\right) - D_{y,u}V_j\left(a, \tilde{T}^p\varphi(b_\nu, x)\right) = 0 \bmod (x)^p.$$

It follows that $D_{a,u}V_j^{p-1}(a, y) - D_{y,u}V_j(a, y) \in \mathcal{R}_{p-1}(a)$ and, by Lemma 5.4,

$$D_{a,u}F_j^{l-1}(a, y) - D_{y,u}F_j(a, y) \in \pi_{l-1}(\mathcal{R}_{p-1}(a)) = \pi_{l-1}(\mathcal{R}_p(a)).$$

On the other hand, $\text{supp}_y [D_{a,u}F_j^{l-1}(a, y) - D_{y,u}F_j(a, y)] \subset \Delta_j$; thus $D_{a,u}F_j^{l-1}(a, y) = D_{y,u}F_j(a, y)$. In virtue of Proposition 3.2, this completes the proof of Lemma 6.5. $\square$

We can now finish the proof of Proposition 6.1. Define $G \in \mathcal{C}^0(A \cup Z)[y]$ by setting $G = F_j$ on Λ_j , for each $j \in J_0$, and $G = 0$ elsewhere. Then $T^l f(b, x) = G(a, \tilde{T}^l \varphi(b, x)) \bmod (x)^{l+1}$, for each $a \in A \cup Z$ and $b \in \varphi^{-1}(a) \cap E$. By Proposition 3.4, the truncation $G^k(a, y)$ is a $\mathcal{C}^k$ Whitney field on $A \cup Z$. $\square$

REFERENCES

1. E. Bierstone, P.D. Milman,, *Composite differentiable functions*, Ann. of Math. (2) **116** ((1982)), 541–558..
2. ———, *Algebras of composite differentiable functions*, Proc. Symp. Pure Math. **40** ((1983)), 127–136..
3. ———, *The Newton diagram of an analytic morphism, and applications to differentiable functions*, Bull. Amer. Math. Soc. **9** ((1983)), 315–318..
4. ———, *Relations among analytic functions, I*, Ann. Inst. Fourier (Grenoble) **37:1** ((1987)), 187–239; and *II*, **37:2** (1987), 49–77..
5. ———, *Semianalytic and subanalytic sets*, Inst. Hautes Études Sci. Publ. Math. **67** ((1988)), 5–42..
6. ———, *Geometric and differential properties of subanalytic sets*, Bull. Amer. Math. Sci. **25** ((1991)), 385–393..

7. E. Bierstone and G.W. Schwarz,, *Continuous linear division and extension of C^∞ -functions*,, Duke Math. J. **50** ((1983)), 233–271..
8. Z. Denkowska, S. Łojasiewicz, J. Stasica,, *Certaines propriétés élémentaires des ensembles sous-analytiques*,, Bull. Polish Acad. Sci. Math. **27** ((1979)), 529–536..
9. Z. Denkowska, S. Łojasiewicz, J. Stasica,, *Sur le théorème du complémentaire pour les ensembles sous-analytiques*,, Bull. Polish Acad. Sci. Math. **27** ((1979)), 537–539..
10. G. Glaeser,, *Fonctions composées différentiables*, Ann. of Math. (2) **77** (1963), 193–209..
11. H. Grauert,, *Über die Deformation isolierter Singularitäten analytischer Mengen*,, Invent. Math. **15** ((1972)), 171–198..
12. R. H. Hardt,, *Stratification of real analytic mappings and images*,, Invent. Math. **28** ((1975)), 193–208..
13. R. H. Hardt,, *Triangulation of subanalytic sets and proper light subanalytic maps*,, Invent. Math. **38** ((1977)), 207–217..
14. H. Hironaka,, *Subanalytic sets*,, in: Number Theory, Algebraic Geometry and Commutative Algebra in Honor of Y. Akizuki, Kinokuniya, Tokyo, (1973), 453–493..
15. H. Hironaka,, *Introduction to the theory of infinitely near singular points*,, Mem. Mat. Inst. Jorge Juan, No. 28, Consejo Superior de Investigaciones Científicas, Madrid, (1974)..
16. S. Łojasiewicz,, *Ensembles semi-analytiques*,, Bures-sur-Yvette, 1964., Inst. Hautes Études Sci.,.
17. S. Łojasiewicz,, *Stratifications et triangulations sous-analytiques*,, Univ. Stud. Bologna Sem. Geom., (1986), 83–89..
18. B. Malgrange,, *Ideals of Differentiable Functions*, Oxford Univ. Press, Bombay, 1966..
19. P.D. Milman,, *The Malgrange-Mather division theorem*,, Topology **16** ((1977)), 395–401..
20. W. Pawłucki,, *On relations among analytic functions and geometry of subanalytic sets*,, Bull. Polish Acad. Sci. Math. **37** ((1989)), 117–125..
21. W. Pawłucki,, *Points de Nash des ensembles sous-analytiques*,, Mem. Amer. Math. Soc. **425** ((1990)).
22. W. Pawłucki,, *On Gabrielov’s regularity condition for analytic mappings*,, Duke Math. J. **65** ((1992)), 299–311..
23. G.W. Schwarz,, *Smooth functions invariant under the action of a compact Lie group*,, Topology **14** ((1975)), 63–68..
24. J. Cl. Tougeron,, *An extension of Whitney’s spectral theorem*,, Inst. Hautes Études Sci. Publ. Math. **40** ((1972)), 139–148..
25. J. Cl. Tougeron,, *Idéaux de Fonctions Différentiables*,, Berlin-Heidelberg-New York, 1972., Springer.,
26. J. Cl. Tougeron,, *Fonctions composées différentiables: cas algébrique*,, Ann. Inst. Fourier (Grenoble) **30** ((1980)), 51–74..
27. H. Whitney,, *Functions differentiable on the boundaries of regions*,, Ann. of Math. **35** ((1934)), 482–485..
28. H. Whitney,, *Differentiable even functions*,, Duke Math. J. **10** ((1943)), 159–160..

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TORONTO, TORONTO, CANADA M5S 1A1.

JAGIELLONIAN UNIVERSITY, INSTITUTE OF MATHEMATICS, UL. REYMONTA 4, 30-059 KRAKÓW, POLAND.